



Pollinator Stewardship Council

1617 White Water Ct. Berthoud, CO 80513

www.pollinatorstewardship.org

April 24, 2020

To: US EPA Office of Pesticide Programs

From: Pollinator Stewardship Council

Re: Docket numbers EPA-HQ-OPP-2012-0329, EPA-HQ-OPP-2011-0865, EPA-HQ-OPP-2011-0920, EPA-HQ-OPP-2008-0844, and EPA-HQ-OPP-2011-0581

The Environmental Protection Agency (EPA) is charged with protecting the health and welfare of both humans and the environment, including pollinators. This letter is written in response to the EPA's Proposed Interim Decision (PID) for acetamiprid, clothianidin/thiamethoxam, dinotefuran and imidacloprid (aka neonicotinoids or neonics). The Proposed Interim Decision maintains the status quo of prophylactic and widespread overuse of these neonicotinoids via foliar, drench and seed treatments within the United States. EPA's statutory regulations must be followed with regard to both the risk assessment process and risk management. Therefore, we assert that EPA has failed to protect both the public and pollinators with the initial conditional registration and has failed to consider new and emerging peer reviewed science which clearly implicates neonics in the decline of pollinators in the U.S. and globally (Potts et al. 2010, Goulson 2019). **We urge EPA to use this review process to impose a total ban on the use of neonics for all outdoor applications, including foliar, drench and seed treatments.** By eliminating the outdoor use of neonics in the U.S., EPA would begin to align the United States with the rest of the world in protecting the public, our ground water, our pollinators and our environment.

As complaints began to mount globally about the harm to pollinators from pesticides, and especially neonics, the first response by EPA and the chemical industry was to convene a Pellston Workshop on testing protocols and risk assessment. This meeting resulted in a book with guidelines that was produced in 2011, but little changed within EPA. In 2012, the EPA called a Science Advisory Panel (SAP) to further evaluate the testing protocols for neonics. In preparation for the SAP panel in 2012, the EPA put together a white paper on pollinators and the issues of concern. EPA stated in section 2.2.1 of the white paper, that the protection goals and associated assessment endpoints for assessing risks to honeybees (*Apis mellifera*) were: "1) protection of pollination services, 2) protection of honey and hive product production and 3) protection of pollinator biodiversity." If these are indeed the accepted protection goals, then EPA needs to stop the overuse of neonics to provide adequate protection for all bees and the pollination services they provide. The latest EPA response has been the establishment of Managed Pollinator Protection Plans (MP3). MP3 guidelines are NOT a substitute for proper label language and enforcement has not and WILL NOT protect pollinators from the resultant harm.

EPA has ignored the best advice from the SAP in that testing of other species beyond *Apis* is needed and that some exposure tests to immature bees should be included. An example of EPA's disregard of that

advice is confirmed by the current statement in the imidacloprid (PID Case 7606) document: “As the pollinator risk assessment framework used by the EPA indicates, honeybees are intended to be reasonable surrogates for other bee species, and conclusions from the weight of evidence for the honeybee can be used to help inform about potential risks to other non-*Apis* species.” This statement is in direct conflict with the SAP recommendations that other species of bees need to be tested and that honey bees are not a sufficient surrogate for all bees. This difference in species response is highlighted by studies that provide direct evidence of negative neonic impact on reproduction in both bumble bees (Whitehorn et al. 2015) and *Osmia*, a solitary bee (Sandrock et al. 2015).

Adding to this evidence is a large multi-year, well-replicated study using neonic seed treatment of canola in Sweden that showed major negative impacts on bumble bees and *Osmia* near treated fields (Rundlöf et al. 2015). While this study failed to show negative impacts on honey bees, it was touted as evidence that little harm is done to *Apis*. These studies indicate that additional bee species should be tested, not ignored, and that honey bees are not an adequate surrogate for other bee species. The reason honey bee colonies did not show evidence of harm is due to the number of individual bees in a colony (20,000-40,000) and this high number allows the colony to have a buffer against pesticide impacts and still show little or no damage even if bees are dying (termed superorganism resilience, Straub et al. 2015).

Honey bee colonies are being negatively impacted by neonics. Recent studies with whole honey bee colonies by Tsvetkov *et al.* (2017) showed that when colonies were near corn crops, bees were exposed to neonics in non-target pollen for 3-4 months which decreased survival. Similarly, Woodcock *et al.* (2017) found that neonicotinoid exposure from nontarget sources, when bees were near canola, reduced overwintering and colony reproduction in both wild bees and honey bees and similar results were found in the UK by Budge *et al.* (2017). Field observations by professional beekeepers confirm these study results.

These field trials, with field-realistic exposure across countries with differing habitat, all show that neonics are leaching into the soils and are then being picked up by non-target plants that bees visit near those agricultural fields. This idea was first demonstrated by the work of Krupke *et al.* (2017) and Botías *et al.* (2015) and, coupled with recent ground water contamination data by neonics (Hladik et al. 2014, Main et al. 2014, Raby et al. 2018), paint a picture of contamination and impact that cannot be ignored. Much of the debate had centered on peer reviewed lab experiments that both the chemical industry and EPA have stated used higher doses than bees would encounter in the field (not field-realistic doses). Indeed, some lab studies may have used a range of dosages that in the upper ranges were not field-realistic. However, this argument is no longer valid given that neonic toxicity increases with time due to bioaccumulation as recently reported in large-scale field tests of chronic exposure or “time-cumulative toxicity” scenarios (Sanchez-Bayo & Tennekes 2020). The EPA uses only exposure dose in determining risk but it must consider the accumulation of neonics in water, soil and plants over time as a more accurate measure of the true risks and the observed negative effects we see on pollinators and other non-target organisms.

The persistence of neonics in the soil (Jones et al. 2014, Krupke et al. 2017) and water (Hladik et al. 2014) make this a mounting problem that will continue for decades even if neonic use stopped today. It will take time for the soil and water to recover. Additionally, the prophylactic use of seed treatments is under-reported (Hitaj 2020). Many invertebrates and vertebrates are being affected by overuse of pesticides as evidenced by the startling studies on the reduction in invertebrate biomass (Hallmann et

al. 2017). We need action now to save bats, birds (Hallmann et al. 2014), butterflies and bees and prevent an “Insect Apocalypse” that threatens the very fabric of our entire ecosystem (Goulson 2019).

Our ground water is also being contaminated with the widespread prophylactic use of neonics on all major crops. The prairie Midwest, where corn and soybean cultivation are dominant, is particularly at risk (Hladik et al. 2014, Main et al. 2014). The Midwest is also the home to over 50% of US honey bee colonies in the summer as they move to this area for honey production. During summer, honey bee colonies take in larger amounts of water to cool the hive and also use water to thin honey and add to food fed to larvae. The bees collect water from the surface of ponds and temporary pools in fields and ditches where the concentrations of pesticides are highest during runoff from fields (Hladik et al. 2014) and where they may not rapidly degrade (Lu et al. 2015). Thus, in addition to the movement of neonics into pollen and nectar via soil contamination; water contamination is also a major source of neonic exposure in honey bee hives.

EPA is bound by the Clean Water and Endangered Species Acts to protect the environment and under FIFRA rules it must take action. We urge EPA to follow its own language in the PIDs which states: *“statutory standard: no unreasonable adverse effects.”* **The scientific evidence is clear: *adverse effects to pollinators and the environment are being seen and documented with the widespread use of neonics.***

FIFRA defines a pesticide as being “misbranded if its labeling bears any statement, graphic representation, or design which is false or misleading in any particular.” Here is a current label for [Admire Pro](#). It is already 67 pages long. This label displays a honey bee in a diamond shape. There is not a red line across the bee that typically accompanies a graphic with a negative message. A user could easily interpret this graphic as meaning it is OK for bees. Further, the current labels on neonics do not contain a caution statement sufficient to protect the health of pollinators and the environment. Under FIFRA rules, if a chemical is misbranded, “the administrator shall remove the registration of the product.” These products do indeed pose unreasonable risk which cannot be mitigated. Neonics are “misbranded” and must be removed from use.

EPA has a mandate to protect the environment. Now is the time for EPA to correct the conditional registrations given in the past and to take into account the mounting scientific evidence that shows a consistent pattern of negative effects on all pollinators at field-realistic exposures that they now regularly encounter. EPA has been subject to the same types of tactics the FDA was subjected to by the tobacco lobby: shoot the messengers and or discredit the science. As with lung cancer victims, beekeepers in the U.S. are suffering because of the inaction of EPA. Economic damage due to the exposure to neonic use is growing (Douglas et al. 2020). Colony losses have gone from 5-10% in the past to well over 50% on an annual basis (Kulhanek et al. 2017). This is unsustainable especially as neonic use is growing (Hitaj et al. 2020). Many beekeeping operations are multi-generational. Beekeepers have not forgotten how to keep bees. The excessive use of neonics in U.S. agriculture has changed not beekeepers’ excellent management practices.

In Europe, a systematic review of neonics resulted in their use being banned (EFSA 2013). European farmers are finding alternatives. Crops are not failing. People are not starving. EPA has tried to respond to pollinator decline with meetings and committees and plans but is still handcuffed by the belief that needed changes will leave farmers vulnerable. We have less toxic alternatives to neonics. Widespread prophylactic treatment flies in the face of responsible agricultural practices. **A total ban on the use of**

neonics in field crops is a necessary first step in protecting the environment we live in and pollinators and the pollination services they provide that feed us daily.

Prepared by: Pollinator Stewardship Council and Drs. Jeff Pettis and Susan Kegley



American Beekeeping Federation

Molly Sausaman
Executive Director
500 Discovery Parkway, Suite 125
Superior, CO 80027
Phone: 720-616-4145
Email: msausaman@abfnet.org
Web: www.abfnet.org



Colorado State Beekeepers Association

Ed Colby, CSBA President
6765 County Road 214
New Castle, CO 81647
president@coloradobeekeepers.org
970-984-0419



Georgia Beekeepers Association

President Linda Tillman
gabepresident@gmail.com
1210 Box Springs Road
Box Springs, GA

404-447-1943



Minnesota
Hobby Beekeepers
Association

Minnesota Hobby Beekeepers Association

Susan Bornstein MHBA President
president@mnbeekeepers.com
11 Lost Rock Lane
St. Paul, MN 55127
651-481-8012



Pollinate Minnesota

Erin Rupp
erin@pollinatemn.org, 612.245.6384
512 2nd Street NE, Mpls, MN 55413



Zen Honeycutt, Founding Executive Director
Mission Viejo, CA. 92691
M. 949-307-6695



Humming for Bees

Jeff Dinsmore

Humming for Bees
jeff@hummingforbees.org
612-720-9084

Jennifer Kraft Leavey, Ph.D.

Principal Academic Professional
Georgia Tech College of Sciences
404-385-7229 jennifer.leavey@cos.gatech.edu

**Metro Atlanta Beekeepers Association**

President Gail Dean

president@metroatlantabeekeepers.org

770-444-2335

562 Rue Montaigne

Stone Mountain, GA 30083

Tri-state Beekeepers Association, Trenton, GA

Eddie Gwaltney, President

407 Moore Road

Lookout Mountain, GA 30750

423-298-7557

Atlanta Natural Beekeeping Meetup

Julia Mahood and Linda Tillman, Founders

4790 Huntley Drive

Atlanta, GA 30342

678-448-2576

The Good Acre

Rhys Williams

Executive Director

612-889-1544

rhys@thegoodacre.org

www.thegoodacre.org

Bird and the Bees Honey, LLC

James Cook

Owner

Monson's Kabekona Bay Hives

Margot Monson, MS Entomology

mpmonson.insx@gmail.com

22 Ludlow Avenue

St. Paul, MN 55108

651-247-5253



Lori D. Cox, Owner/Operator
Roots Return Heritage Farm, LLC
 14525 County Road 40 Carver, MN 55315
 425-241-2515

Muskrat Farm

Keith Chrysler & Laurie Schneider, farmers and beekeepers
 9503 Norell Avenue
 Stillwater, MN 55082



Tara Beekeepers Association
 752 Main Street, Forest Park, GA 30297
<https://www.tarabeekeepers.org/>
 tarabeekeepers@gmail.com
 404-369-3809

California Minnesota Honey Farms

Jeff Anderson
 721 Wells Street
 Eagle Bend MN 56446



Honey Pond Farm
 Jennifer Berry

747 Madison St.
Comer, GA
30629
jennifer@honeypondfarm.com

MtnHoney

Virginia Webb
349 Gastley Road
Clarksville, GA 30523
706-754-7062

Niwot Honey Farm


P.O. Box 33, Niwot, Colorado 80544
Niwot, Colorado
Tom Theobald Beekeeper/Owner

**Tri-County Beekeepers**

Bobby Chaisson
President
4664 Highway 51 South
Lula, Georgia 30554

**Adee Honey Farms**

515 Jay Street
Bruce, SD 57220
605-627-5621



Bee Squared Apiaries

1617 White Water Ct.
 Berthoud, CO 80513
 beth@bethsbees.com
<http://bethsbees.com>

Bird AND THE Bees

Bird and the Bees Honey, LLC

N9453 Pleasant Hill Rd.
 Iola, WI 54945
 birdandthebeeshoney@gmail.com
 James Cook

Cox Honey of Utah

Darren Cox
 PO Box 435
 Menden, Utah 84323



Hackenberg Apiaries
 Lewisburg, Pennsylvania
 570-568-2337
 hackenberghoney@gmail.com
<https://hackenbergapiaries.us/>

Wilson's Honey

6626 Stitt Rd, Altamont, NY 12009
stephenwilson282@gmail.com



Eastern Piedmont Beekeepers Association

431 Hale Ln
Athens GA 30607
Dan Long, President



Mac's Backyard Bees

Bob McDonell
350 Sherwood Drive
Carol Stream, IL 60188
beekeeperbob@gmail.com
630-940-7655

James Doan

PO Box 25
West Clarksville, NY
14786

References:

Botías et al. 2015. Neonicotinoid residues in wildflowers, a potential route of chronic exposure for bees. *Environ. Sci. Technol.* **49**, 12731–12740.

Budge et al. 2015. Evidence for pollinator cost and farming benefits of neonicotinoid seed coatings on oilseed rape. *Scientific Reports.* **5**: 12574.

Douglas et al. 2020. County-level analysis reveals a rapidly shifting landscape of insecticide hazard to honey bees (*Apis mellifera*) on US farmland. *Scientific Reports.* **10**: 797.

European Food Safety Authority (EFSA) 2013. Conclusion on the peer review of the pesticide risk assessment for bees for the active substance thiamethoxam. *EFSA J.* **11**, 3066. doi:10.2903/j.efsa.2013.3066

EFSA 2013. Conclusion on the peer review of the pesticide risk assessment for bees for the active substance clothianidin. *EFSA J.* **11**, 3067. doi:10.2903/j.efsa.2013.3067

EFSA 2013. Conclusion on the peer review of the pesticide risk assessment for bees for the active substance imidacloprid. *EFSA J.* **11**, 3068. doi:10.2903/j.efsa.2013.3068

Goulson, D. 2019. The insect apocalypse, and why it matters. *Current Biology.* **29** (19)967-971.

Hallmann et al. 2014 Declines in insectivorous birds are associated with high neonicotinoid concentrations. *Nature.* **511**: 341–343.

Hallmann et al. 2017. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. October 18, 2017 *PLoS One* 0185809

Henry et al. 2012. A common pesticide decreases foraging success and survival in honey bees. *Science* **336**, 348–350.

Hitaj et al. 2020. Sowing Uncertainty: What We Do and Don't Know about the Planting of Pesticide-Treated Seed. *BioScience.* doi.org/10.1039/biosci

Hladik et al. 2014. Widespread occurrence of neonicotinoid insecticides in streams in a high corn and soybean producing region, USA. *Environmental Pollution* **193**: 189-196.

Hladik et al. 2018. Environmental Risks and Challenges Associated with Neonicotinoid Insecticides. *Environ. Sci. Technol.* **52**, 6, 3329.

Jones et al. 2014. Neonicotinoid concentrations in arable soils after seed treatment applications in preceding years. *Pest Manag. Sci.* **70**, 1780–1784.

Krupke et al. 2017. Planting of neonicotinoid-treated maize poses risks for honey bees and other non-target organisms over a wide area without consistent crop yield benefit. *Journal of Applied Ecology.* **54**: 1449–1458. doi: 10.1111/1365-2664.12924

Kulhanek et al. 2017. A national survey of managed honey bee 2015–2016 annual colony losses in the USA. *J. of Apicultural Research* **56**: 328-340.

Lu et al. 2015. Quantum Yields for Direct Photolysis of Neonicotinoid Insecticides in Water: Implications for Exposure to Nontarget Aquatic Organisms. *Environ. Sci. Technol. Lett.* **2**, 7, 188-192.

Main et al. 2014. Widespread Use and Frequent Detection of Neonicotinoid Insecticides in Wetlands of Canada's Prairie Pothole Region. *PLoS ONE* **9**(3): e92821. doi:10.1371/journal.pone.0092821

Potts et al. 2010. Global pollinator declines: Trends, impacts and drivers. *Trends Ecol. Evol.* **25**, 345–353.

Raby et al. 2018. Acute toxicity of 6 neonicotinoid insecticides to freshwater invertebrates. *Environmental Toxicology and Chemistry.* **37**(5): 1430-1445.

Rundlöf et al. 2015. Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature.* **521**: 77–80.

Sandrock et al. 2014. Sublethal neonicotinoid insecticide exposure reduces solitary bee reproductive success. *Agric. For. Entomol.* **16**: 119–128.

Sanchez-Bayo and Tennekes. 2020. Time-cumulative toxicity of neonicotinoids: experimental evidence and implications for environmental risk assessments. *International J. of Environmental Research and Public Health.* 17: 1629.

Straub et al. 2015. Superorganism resilience: eusociality and susceptibility of ecosystem service providing insects to stressors. *Current Opinion in Insect Science.* Vol 12: 109-112.

Tsvetkov et al. 2017. Chronic exposure to neonicotinoids reduces honey bee health near corn crops *Science.* Vol. 356, Issue **6345**: 1395-1397.

Whitehorn et al. 2012. Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* Vol. **336**: 351–352.

Woodcock et al. 2017. Country-specific effects of neonicotinoid pesticides on honey bees and wild bees. *Science* Vol. **356**: 1393-1395.